

Squeezing lepton pairs out of broken symmetries

A. K. Dutt-Mazumder*, C. Gale*, A. Majumder* and O. Teodorescu*

**Department of Physics, McGill University, Montreal, QC, Canada H3A 2T8*

Abstract. We discuss two possible signatures of symmetry breaking that can appear in dilepton spectra, as measured in relativistic heavy ion collisions. The first involves scalar-vector meson mixing and is related to the breaking of Lorentz symmetry by a hot medium. The second is related to the breaking of Furry's theorem by a charged quark-gluon plasma. Those signals will be accessible to upcoming measurements to be performed at the GSI, RHIC, and the LHC.

INTRODUCTION

Electromagnetic radiation, especially lepton pairs, represents probes of choice for the study of heavy ion reactions. Owing to the smallness of their final state interaction they travel essentially unscathed from creation to detection. Furthermore, via Vector Meson Dominance (VMD) [1], real and virtual photons are ideal as signatures of the in-medium behaviour of vector mesons [2]. This aspect has been highlighted in a theoretical interpretation [3] of the low dilepton invariant mass measurements performed by the CERES collaboration [4]. In short, it appears that the behaviour of hot and dense strongly interacting matter can indeed be revealed by electromagnetic observables [5].

In this work we shall outline how symmetry breaking effects can manifest themselves in dilepton spectra. The first of those involve the possibility of $\rho - a_0$ mixing in dense nuclear matter. The size of this contribution is density-dependent and goes to zero in vacuum, on account of Lorentz symmetry. We show that such a mixing opens up a new channel in dilepton production and will modify the spectrum in the ϕ invariant mass region. Secondly, we show that Furry's theorem is no longer obeyed in a baryon rich quark gluon plasma. This has the consequence of permitting new processes like $gg \rightarrow l^+ l^-$. We evaluate this contribution and compare it with $q\bar{q} \rightarrow l^+ l^-$. We will consider only dielectrons here and the details of the calculations will only be sketched; the interested reader is invited to consult the references quoted in the text. The observables we discuss in this work are within reach of measurements soon to be performed by the HADES collaboration at the GSI, and by the PHENIX collaboration at RHIC.

DILEPTONS FROM MESON MIXING IN DENSE HADRONIC MATTER

There exists physical processes which are forbidden in free space but can take place in matter. Those are related to medium-induced symmetry breaking effects. The interaction respects the symmetry, but it is broken by the ground state. For instance, in matter, Lorentz symmetry is lost which leads to the mixing of different spin states even when the interaction Lagrangian respects all the required symmetry properties. A well-known example of this is σ - ω mixing in nuclear matter as discussed by Chin [6]. Another manifestation lies in the fact that different polarization states of vector fields can have different dispersion relations [2].

Our goal here is to identify the effects of scalar and vector meson mixing on dilepton production rates, which could be observed in high energy heavy ion experiments. The mixing of mesons considered here include both isoscalar and isovector channels. To be more specific, we estimate the rate of dilepton production from σ - ω and ρ - a_0 mixing. Some attention was paid to the former in the context of heavy-ion collisions [7, 8], and the importance of the later was shown recently [9].

In this first part we present thermal production rates of dileptons induced by scalar-vector mixing in the isoscalar and isovector channels. Those results are compared against π - π , K - $\bar{K}$ annihilation contributions, which are treated here as standard candles. Note that processes involving mixing of different G-parity states like σ - ω or ρ - a_0 are allowed only in matter which is not invariant under charge conjugation. Therefore, $\pi\pi \rightarrow e^+ e^-$ through the coupling to nucleons via σ and ω , or $\pi\eta \rightarrow e^+ e^-$ mediated by ρ - a_0 mixing, can take place in matter with a finite nucleon chemical potential.

Hence it is natural to look for such density-driven effects in experiments involving relatively low energy collisions where one expects to have a relatively large chemical potential.

The interaction Lagrangian used in the present model can be written as

$$\mathcal{L}_{int} = g_\sigma \bar{\psi} \phi_\sigma \psi + g_{a_0} \bar{\psi} \phi_{a_0, \alpha} \tau^\alpha \psi + g_{\omega NN} \bar{\psi} \gamma_\mu \psi \omega^\mu + \quad (1)$$

$$g_\rho [\bar{\psi} \gamma_\mu \tau^\alpha \psi + \frac{\kappa_\rho}{2m_n} \bar{\psi} \sigma_{\mu\nu} \tau^\alpha \partial^\nu] \rho^\mu_\alpha, \quad (2)$$

where ψ , ϕ_σ , ϕ_{a_0} , ρ and ω correspond to nucleon, σ , a_0 , ρ and ω fields, and the τ 's are Pauli matrices. From this point on we use s and v to denote scalar and vector mesons, *i.e.* $s = \sigma, a_0$ and $v = \omega, \rho$. It is understood that $\kappa_\omega = 0.0$.

The polarization vector through which the scalar meson couples to the vector meson via the n - n loop is given by

$$\Pi_\mu(q_0, |\vec{q}|) = 2ig_s g_v \int \frac{d^4 k}{(2\pi)^4} \text{Tr}[G(k) \Gamma_\mu G(k+q)]. \quad (3)$$

where 2 is an isospin factor and the vertex for v - nn coupling is:

$$\Gamma_\mu = \gamma_\mu - \frac{\kappa_v}{2m_n} \sigma_{\mu\nu} q^\nu. \quad (4)$$

$G(k)$ is the in-medium nucleon propagator [10].

ρ - a_0 mixing opens up a new channel for dilepton production in dense and hot nuclear matter: $\pi + \eta \rightarrow e^+ + e^-$, accessible through n - n excitations. The Feynman diagram for this process is shown in Fig. 1. Similarly, the contribution for $\pi + \pi \rightarrow \sigma \rightarrow \omega \rightarrow e^+ + e^-$ can be evaluated from a diagram obtained from that in Fig. 1 by replacing one of the incoming π 's by η and by performing the following substitutions: $a_0 \rightarrow \sigma$ and $\rho \rightarrow \omega$.




FIGURE 1. The Feynman diagram for the process $\pi + \eta \rightarrow e^+ + e^-$.

Dilepton rates

Once we obtain the relevant cross-section for the dilepton production channel, the thermal production rate of the lepton pairs (number of reactions per unit time, per unit volume $R_{12}^{e^+e^-} = dN_{e^+e^- \text{ pairs}}/d^4x$) can be estimated in the independent particle approximation of relativistic kinetic theory:

$$\frac{dR_{12}^{e^+e^-}}{dM^2} = \int \frac{d^3 k_1}{(2\pi)^3} f_1(\mathbf{k}_1) \int \frac{d^3 k_2}{(2\pi)^3} f_2(\mathbf{k}_2) \frac{d\sigma_{12}^{e^+e^-}(s, M^2)}{dM^2} v_{rel} \quad (5)$$

where f_1, f_2 are the thermal distributions of the 1,2 species and $v_{rel} = \frac{\lambda^{1/2}(s, m_1^2, m_2^2) \sqrt{s}}{2E_1 E_2}$ with $\lambda(x, y, z) = x^2 + y^2 + z^2 - 2xy - 2xz - 2yz$ is the triangle function.

Fig. 2 shows the total dilepton yield with the combined effect of mixing in the isoscalar and isovector channels. We have chosen conditions that are representative of collisions to occur at the GSI, and we show the pion and kaon annihilation channels separately. It is important to see that rates induced by the mixing can be significant in certain windows of density and temperature. Particularly, mixing induces a higher yield of dileptons in the invariant mass region between the ρ and ϕ peaks, and is strongly density dependent. The mixing in the isoscalar channel contribute essentially at the ω peak ¹. On the other hand, the isovector channel seems to provide a better probe as it contributes

¹ Technically, the $\pi\pi$ annihilation channel should treat the s - and the p -wave annihilation coherently. The interference effects have been found to be negligible.




FIGURE 2. Rates at $T = 75$ MeV with mixing in both isoscalar and isovector channels.

in the tail region of the two pion annihilation contribution. It should be stressed here that broadening or dropping ρ meson mass would even favour our observation. In that case the π - π background would be pushed more towards lower invariant mass region bringing the a_0 peak into a clearer relief. Importantly, the upcoming HADES experiment [12] has the necessary resolution, and will operate in the appropriate energy regime to adequately investigate the physics discussed here.

DILEPTON EMISSION FROM A BARYON-RICH QUARK GLUON PLASMA

At zero temperature, diagrams in QED that contain a fermion loop with an odd number of photon vertices are cancelled by an equal and opposite contribution coming from the same diagram with fermion lines running in the opposite direction (Furry's theorem [13]). This statement can also be generalized to QCD for processes with two gluons and an odd number of photon vertices.

At finite temperature and density, the corresponding quantity to consider in connection with Furry's theorem is $\sum_n \langle n | A_{\mu_1} A_{\mu_2} \dots A_{\mu_{2n+1}} | n \rangle e^{-\beta(E_n - \mu Q_n)}$, where $\beta = 1/T$, μ is a chemical potential, and A_μ is an electromagnetic gauge field operator. Here, the effect of the charge conjugation operator C is such that $C|n\rangle = e^{i\phi}|-n\rangle$, where $|-n\rangle$ is a state in the ensemble with the same number of antiparticles as there are particles in $|n\rangle$ and vice-versa. If $\mu = 0$, upon multiple insertions of the unit operator $C^{-1}C$ we get

$$\langle n | A_{\mu_1} A_{\mu_2} \dots A_{\mu_{2n+1}} | n \rangle e^{-\beta E_n} = - \langle -n | A_{\mu_1} A_{\mu_2} \dots A_{\mu_{2n+1}} | -n \rangle e^{-\beta E_n}. \quad (6)$$

The sum over all states will contain the mirror term $\langle -n | A_{\mu_1} A_{\mu_2} \dots A_{\mu_{2n+1}} | -n \rangle e^{-\beta E_n}$ with the same thermal weight. Therefore

$$\sum_n \langle n | A_{\mu_1} A_{\mu_2} \dots A_{\mu_{2n+1}} | n \rangle e^{-\beta E_n} = 0 \quad (7)$$

and Furry's theorem still holds.

However, if $\mu \neq 0$ (unequal number of particles and antiparticles), then

$$\langle n | A_{\mu_1} A_{\mu_2} \dots A_{\mu_{2n+1}} | n \rangle e^{-\beta(E_n - \mu Q_n)} = - \langle -n | A_{\mu_1} A_{\mu_2} \dots A_{\mu_{2n+1}} | -n \rangle e^{-\beta(E_n - \mu Q_n)}. \quad (8)$$

The mirror term this time is $\langle -n | A_{\mu_1} A_{\mu_2} \dots A_{\mu_{2n+1}} | -n \rangle e^{-\beta(E_n + \mu Q_n)}$, with a different thermal weight. Thus

$$\sum_n \langle n | A_{\mu_1} A_{\mu_2} \dots A_{\mu_{2n+1}} | n \rangle e^{-\beta(E_n - \mu Q_n)} \neq 0. \quad (9)$$

One may say that the medium, being charged, manifestly breaks charge conjugation invariance and these Green's functions are thus finite and will lead to the appearance of new processes in a perturbative expansion. Again, the appearance of processes that can be related to symmetry breaking in a medium has been noted before [6, 14].




FIGURE 3. The two gluon photon effective vertex as the sum of two diagrams with quark number running in opposite directions.

Let us now focus our attention on the diagrams of Fig. 3 for the case of two gluons and a photon attached to a quark loop. Such a process does not exist at zero temperature or even at finite temperature and zero density. At finite density this leads to a new source of dileptons: $gg \rightarrow e^+e^-$. In order to obtain the full matrix element one must coherently sum contributions from both diagrams which have fermion number running in opposite directions. The amplitudes for $T^{\mu\rho\nu} (= T^{\mu\rho\nu} + T^{\nu\rho\mu})$ are

$$T^{\mu\rho\nu} = \frac{1}{\beta} \sum_{n=-\infty}^{\infty} \int_{-\infty}^{\infty} eg^2 \text{tr}[t^a t^b] \frac{d^3 q}{(2\pi)^3} \text{Tr}[\gamma^\mu \gamma^\beta \gamma^\rho \gamma^\delta \gamma^\nu \gamma^\alpha] \frac{(q+p-k)_\alpha q_\beta (q+p)_\delta}{(q+p-k)^2 q^2 (q+p)^2},$$

$$T^{\nu\rho\mu} = \frac{1}{\beta} \sum_{n=-\infty}^{\infty} \int_{-\infty}^{\infty} eg^2 \text{tr}[t^a t^b] \frac{d^3 q}{(2\pi)^3} \text{Tr}[\gamma^\nu \gamma^\delta \gamma^\rho \gamma^\beta \gamma^\mu \gamma^\alpha] \frac{(q+k-p)_\alpha q_\beta (q-p)_\delta}{(q+k-p)^2 q^2 (q-p)^2}, \quad (10)$$

and they involve a trace over colour matrices, and a sum over Matsubara frequencies.

Again note that the extension of Furry's theorem to finite temperature does not hold at finite density: if we set $n \rightarrow -n-1$ we note that $q_0 \not\rightarrow -q_0$ and as a result $T^{\mu\rho\nu}(\mu, T) \neq -T^{\nu\rho\mu}(\mu, T)$. Of course, If we now let the chemical potential go to zero ($\mu \rightarrow 0$) we note that for the transformation $n \rightarrow -n-1$ we obtain $q_0 \rightarrow -q_0$ and thus $T^{\mu\rho\nu}(0, T) \rightarrow -T^{\nu\rho\mu}(0, T)$. The analysis for fermion loops with larger number of vertices is essentially the same.

Dilepton rates

To calculate the contribution made by the diagram of Fig. 3 to the dilepton spectrum emanating from a quark gluon plasma we calculate the imaginary part of the photon self-energy containing the above diagram as an effective vertex (see [11] for details). We calculate in the limit of photon three momentum $\vec{p} = 0$. The imaginary part of the considered self energy contains various cuts. We concentrate solely on the cut that represents the process of gluon-gluon to e^+e^- . This is a new process which to our knowledge has not been discussed before. The other possible cuts represent essentially finite density contributions to other known processes of dilepton production. The differential production rate for pairs of massless leptons with total energy E and and total momentum $\vec{p} = 0$ is given in terms of the discontinuity in the photon self-energy [2]. We use the $q\bar{q} \rightarrow e^+e^-$ Born process as a baseline.

The initial temperatures of the plasma formed at RHIC and LHC have been estimated to lie in the range from 300-800 MeV [15, 16]. For this exploratory calculation we use a conservative estimate of $T = 400$ MeV. To evaluate the effect of a finite chemical potential we perform the calculation with two extreme values of chemical potential $\mu = 0.1T$ (1st plot in Fig. 4) and $\mu = 0.5T$ (2nd plot Fig. 4) [17]. The calculation, is performed for three massless flavours of quarks. The differential rate for the production of dileptons with an invariant mass from 0.5 to 2.5 GeV is presented. On purpose, we avoid regions where the gluons become very soft. In the plots, the dashed line is the rate from tree level $q\bar{q}$; the solid line is that from the process $gg \rightarrow e^+e^-$. We note that in both cases the gluon-gluon process dominates at low energy and dies out at higher energy leaving the $q\bar{q}$ process dominant. Importantly, the crossing point lies near the continuum between the ϕ and the J/ψ , raising an optimistic prospect for the successful detection of the new contribution.

SUMMARY

We have considered the lepton pair signature of symmetry breaking effects in two different energy regimes. Those effects are present in finite density media that break charge conjugation invariance. They vanish identically in vacuum,




FIGURE 4. The differential production rate of back to back dileptons from two processes. Invariant mass runs from 0.5 GeV to 2.5 GeV. The dashed line represents the contribution from the process $q\bar{q} \rightarrow e^+e^-$. The solid line corresponds to the process $g\bar{g} \rightarrow e^+e^-$. Temperature is 400 MeV. Quark chemical potential is 0.1T. The second figure is the same as the first but with $\mu = 0.5T$

are therefore genuine in-medium manifestations, and are a testimony to the richness of the many-body problem. Theoretical refinements are needed before truly quantitative predictions can be made. In the lower energy sector, many-body theory has to be introduced in order to consistently extend the current vector meson spectral densities to include meson mixing effects [18]. At higher energies, we have begun a study of the effects of Hard Thermal Loop resummation [19, 20]. Finally, our rates will be inserted in appropriate dynamical models. As those works are in progress, it is stimulating to note that the signals discussed in this work are all accessible to the new generation of lepton pair measurements.

ACKNOWLEDGMENTS

This work was supported in part by the Natural Sciences and Engineering Research Council of Canada, and in part by the Fonds FCAR of the Quebec government.

REFERENCES

1. J. J. Sakurai, *Currents and Mesons*, University of Chicago Press, Chicago, 1969.
2. C. Gale and J. I. Kapusta, *Nucl. Phys.*, **B357**, 65 (1991).
3. R. Rapp and J. Wambach, *Adv. Nucl. Phys.*, **25**, 1 (2000), and references therein.
4. See H. Appelshaeuser, Proceedings of Quark Matter 2001, *Nucl. Phys. A*, in press, and references therein.
5. See, for example, Charles Gale, Proceedings of Quark Matter 2001, *Nucl. Phys. A*, in press, and references therein.
6. S. A. Chin, *Ann. Phys.*, **108**, 301, (1977)
7. G. Wolf, B. Friman, and M. Soyeur, *Nucl. Phys.*, **A640**, 129, (1998).
8. K. Saito, K. Tsushima, A. W. Thomas, and A. G. Williams, *Phys. Lett.*, **B433**, 243 (1998).
9. O. Teodorescu, A.K. Dutt-Mazumder, and C. Gale, *Phys.Rev. C*, **61**, 051901 (2000); *Phys. Rev. C*, **63**, 034903 (2001).
10. B. D. Serot and J. D. Walecka, *Adv. Nucl. Phys.*, **16**, 1 (1986).
11. A. Majumder and C. Gale, *Phys. Rev. D*, **63**, 114008 (2001), and erratum in press.
12. J. Friese *et al.*, *Prog. Part. Nucl. Phys.*, **42**, 235 (1999).
13. W. H. Furry, *Phys. Rev.*, **51**, 125 (1937).
14. H. A. Weldon, *Phys. Lett. B*, **274**, 133 (1992).
15. X. N. Wang, *Phys. Rept.*, **280**, 287 (1997).
16. R. Rapp, hep-ph/0010101.
17. K. Geiger and J. I. Kapusta, *Phys. Rev. D*, **47**, 4905 (1993); N. George, Proceedings of Quark Matter 2001, *Nucl. Phys. A*, in press.
18. O. Teodorescu, A. K. Dutt-Mazumder, and C. Gale, in preparation.
19. E. Braaten and R. D. Pisarski, *Nucl. Phys.*, **B337**, 569 (1990).
20. A. Majumder and C. Gale, *Proceedings of MRST 2001*, to appear, and in preparation.